

# The complemented subspace problem in Banach lattices: A counterexample

David de Hevia

work **in progress** with Gonzalo Martínez Cervantes, Alberto Salguero, and Pedro Tradacete

Instituto de Ciencias Matemáticas



## Positivity XI

Ljubljana, July 10, 2023

Supported by grants CEX2019-000904-S and PID2020-116398GB-I00 funded by MCIN/AEI/10.13039/501100011033 and by an FPU grant FPU20/03334 funded by Ministerio de Universidades.

# Table of Contents

- 1 A general overview of the question
- 2 Some remarks about  $PS_2$
- 3 Concluding remarks

# The Complemented Subspace Problem

*One of the most important problems in the theory of Banach lattices, which is still open, is whether any complemented subspace of a Banach lattice must be linearly isomorphic to a Banach lattice.*

P. Casazza, N. Kalton and L. Tzafriri (1987)

# The Complemented Subspace Problem

*One of the most important problems in the theory of Banach lattices, which is still open, is whether any complemented subspace of a Banach lattice must be linearly isomorphic to a Banach lattice.*

P. Casazza, N. Kalton and L. Tzafriri (1987)

**Definition.** A **Banach lattice** is a real Banach space  $(X, \|\cdot\|)$  equipped with a lattice order  $\leq$  which is compatible with the linear structure of  $X$  (1) and with its norm (2) in the sense that

- 1 If  $x \leq y$ , then  $x + z \leq y + z$  and  $ax \leq ay$  for any  $a \in \mathbb{R}^+$
- 2 If  $|x| \leq |y|$ , then  $\|x\| \leq \|y\|$ .

# The Complemented Subspace Problem

*One of the most important problems in the theory of Banach lattices, which is still open, is whether any complemented subspace of a Banach lattice must be linearly isomorphic to a Banach lattice.*

P. Casazza, N. Kalton and L. Tzafriri (1987)

**Definition.** A **Banach lattice** is a real Banach space  $(X, \|\cdot\|)$  equipped with a lattice order  $\leq$  which is compatible with the linear structure of  $X$  (1) and with its norm (2) in the sense that

- ① If  $x \leq y$ , then  $x + z \leq y + z$  and  $ax \leq ay$  for any  $a \in \mathbb{R}^+$
- ② If  $|x| \leq |y|$ , then  $\|x\| \leq \|y\|$ .

**CSP in Banach lattices:** Are complemented subspaces in Banach lattices isomorphic to Banach lattices?

# The Complemented Subspace Problem

*One of the most important problems in the theory of Banach lattices, which is still open, is whether any complemented subspace of a Banach lattice must be linearly isomorphic to a Banach lattice.*

P. Casazza, N. Kalton and L. Tzafriri (1987)

**Definition.** A **Banach lattice** is a real Banach space  $(X, \|\cdot\|)$  equipped with a lattice order  $\leq$  which is compatible with the linear structure of  $X$  (1) and with its norm (2) in the sense that

- ① If  $x \leq y$ , then  $x + z \leq y + z$  and  $ax \leq ay$  for any  $a \in \mathbb{R}^+$
- ② If  $|x| \leq |y|$ , then  $\|x\| \leq \|y\|$ .

**CSP in Banach lattices:** Are complemented subspaces in Banach lattices isomorphic to Banach lattices?

**Terminology:** By an **isomorphism**  $T : E \rightarrow F$  ( $E, F$  Banach spaces) we mean a bijective continuous linear mapping such that  $T^{-1}$  is also continuous.

We say that a subspace  $F$  of a Banach space  $E$  is **complemented** if there exists a continuous linear mapping  $P : E \rightarrow E$ , with  $P \circ P = P$ , such that  $P(E) = F$ .

# The Complemented Subspace Problem

Let  $X$  be a Banach lattice and let  $E \subset X$  be a Banach space complemented by a linear projection  $P : X \rightarrow X$ . An important **remark**:

We are **not assuming** any relation between the lattice structure of  $X$  and the projection  $P$ .

# The Complemented Subspace Problem

Let  $X$  be a Banach lattice and let  $E \subset X$  be a Banach space complemented by a linear projection  $P : X \rightarrow X$ . An important **remark**:

We are **not assuming** any relation between the lattice structure of  $X$  and the projection  $P$ .

- If  $P$  is a **lattice homomorphism** (that is,  $P(x \vee y) = Px \vee Py$ ), then  $E$  is a sublattice of  $X$ .



# The Complemented Subspace Problem

Let  $X$  be a Banach lattice and let  $E \subset X$  be a Banach space complemented by a linear projection  $P : X \rightarrow X$ . An important **remark**:

We are **not assuming** any relation between the lattice structure of  $X$  and the projection  $P$ .

- If  $P$  is a **lattice homomorphism** (that is,  $P(x \vee y) = Px \vee Py$ ), then  $E$  is a sublattice of  $X$ .
- If  $P$  is **positive** (that is,  $Px \geq 0$  whenever  $x \geq 0$ ), then  $E$  with the order inherited, its lattice operations given by

$$x \vee_E y = P(x \vee y), \quad x \wedge_E y = P(x \wedge y) \quad \text{and} \quad |x|_E = P(|x|),$$

and with the renorming  $\|x\| = \|P|x|\|$  (for  $x \in E$ ) is a Banach lattice.

# Some Banach spaces which are not Banach lattices

Let  $X$  be a Banach lattice.

**Criteria**

**Examples**

# Some Banach spaces which are not Banach lattices

Let  $X$  be a Banach lattice.

## Criteria

- $c_0 \not\subset X$  and  $\ell_1 \not\subset X \iff X$  reflexive.

## Examples

James  $\mathcal{J}$ .

# Some Banach spaces which are not Banach lattices

Let  $X$  be a Banach lattice.

## Criteria

- $c_0 \not\subset X$  and  $\ell_1 \not\subset X \iff X$  reflexive.
- $X$  has an unconditional basic sequence.

## Examples

James  $\mathcal{J}$ .

Gowers-Maurey (1993).

# Some Banach spaces which are not Banach lattices

Let  $X$  be a Banach lattice.

## Criteria

- $c_0 \not\subset X$  and  $\ell_1 \not\subset X \iff X$  reflexive.
- $X$  has an unconditional basic sequence.
- $X$  has the *GL-lust* property.

## Examples

James  $\mathcal{J}$ .

Gowers-Maurey (1993).

$\mathcal{H}^\infty(\mathbb{D})$ ,  $Z_2$  (Kalton-Peck)...

# Some Banach spaces which are not Banach lattices

Let  $X$  be a Banach lattice.

## Criteria

- $c_0 \not\subset X$  and  $\ell_1 \not\subset X \iff X$  reflexive.
- $X$  has an unconditional basic sequence.
- $X$  has the *GL-lust* property.

## Examples

James  $\mathcal{J}$ .

Gowers-Maurey (1993).

$\mathcal{H}^\infty(\mathbb{D})$ ,  $Z_2$  (Kalton-Peck)...

**Obstacle:** These criteria remain valid when  $X$  is a complemented subspace of a Banach lattice.

# Some Banach spaces which are not Banach lattices

Let  $X$  be a Banach lattice.

## Criteria

- $c_0 \not\subset X$  and  $\ell_1 \not\subset X \iff X$  reflexive.
- $X$  has an unconditional basic sequence.
- $X$  has the *GL-lust* property.

## Examples

James  $\mathcal{J}$ .

Gowers-Maurey (1993).

$\mathcal{H}^\infty(\mathbb{D})$ ,  $Z_2$  (Kalton-Peck)...

**Obstacle:** These criteria remain valid when  $X$  is a complemented subspace of a Banach lattice.

Free Banach lattices provide a (not very tractable) criterion to distinguish between the two:

# Some Banach spaces which are not Banach lattices

Let  $X$  be a Banach lattice.

## Criteria

- $c_0 \not\subset X$  and  $\ell_1 \not\subset X \iff X$  reflexive.
- $X$  has an unconditional basic sequence.
- $X$  has the *GL-lust* property.

## Examples

James  $\mathcal{J}$ .

Gowers-Maurey (1993).

$\mathcal{H}^\infty(\mathbb{D})$ ,  $Z_2$  (Kalton-Peck)...

**Obstacle:** These criteria remain valid when  $X$  is a complemented subspace of a Banach lattice.

Free Banach lattices provide a (not very tractable) criterion to distinguish between the two:

- $X$  is complemented in a Banach lattice  $\iff X$  is complemented in  $\text{FBL}[X]$ .



# Some Banach spaces which are not Banach lattices

Let  $X$  be a Banach lattice.

## Criteria

- $c_0 \not\subset X$  and  $\ell_1 \not\subset X \iff X$  reflexive.
- $X$  has an unconditional basic sequence.
- $X$  has the *GL-lust* property.

## Examples

James  $\mathcal{J}$ .

Gowers-Maurey (1993).

$\mathcal{H}^\infty(\mathbb{D})$ ,  $Z_2$  (Kalton-Peck)...

**Obstacle:** These criteria remain valid when  $X$  is a complemented subspace of a Banach lattice.

Free Banach lattices provide a (not very tractable) criterion to distinguish between the two:

- $X$  is complemented in a Banach lattice  $\iff X$  is complemented in  $\text{FBL}[X]$ .
- $X$  is isomorphic to a Banach lattice  $\iff$  there is an ideal  $I \subset \text{FBL}[X]$  such that  $\text{FBL}[X] = I \oplus X$ .

# Some answers and some open questions

## Positive answers

- Every **1-complemented** subspace of an  $L_p$ -space ( $1 \leq p < \infty$ ) is an  $L_p$ -space (Bernau-Lacey 1974).

## Conjectures (?)

- Every complemented subspace of  $L_1[0, 1]$  is isomorphic to  $\ell_1$  or  $L_1[0, 1]$ .

# Some answers and some open questions

## Positive answers

- Every **1-complemented** subspace of an  $L_p$ -space ( $1 \leq p < \infty$ ) is an  $L_p$ -space (Bernau-Lacey 1974).
- Every **1-complemented** subspace of a **separable**  $C(K)$ -space is isomorphic to a  $C(K)$ -space (Benyamini 1973).

## Conjectures (?)

- Every complemented subspace of  $L_1[0, 1]$  is isomorphic to  $\ell_1$  or  $L_1[0, 1]$ .
- Every complemented subspace of  $C[0, 1]$  is isomorphic to a  $C(K)$ -space.

# Some answers and some open questions

## Positive answers

- Every **1-complemented** subspace of an  $L_p$ -space ( $1 \leq p < \infty$ ) is an  $L_p$ -space (Bernau-Lacey 1974).
- Every **1-complemented** subspace of a **separable**  $\mathcal{C}(K)$ -space is isomorphic to a  $\mathcal{C}(K)$ -space (Benyamini 1973).
- In the **complex case**: every **1-complemented** subspace of a space with 1-unconditional basis also has 1-unconditional basis. (Kalton-Wood 1976)

## Conjectures (?)

- Every complemented subspace of  $L_1[0, 1]$  is isomorphic to  $\ell_1$  or  $L_1[0, 1]$ .
- Every complemented subspace of  $C[0, 1]$  is isomorphic to a  $C(K)$ -space.
- Every complemented subspace of a space with unconditional basis has unconditional basis.

# Some answers and some open questions

## Positive answers

- Every **1-complemented** subspace of an  $L_p$ -space ( $1 \leq p < \infty$ ) is an  $L_p$ -space (Bernau-Lacey 1974).
- Every **1-complemented** subspace of a **separable**  $C(K)$ -space is isomorphic to a  $C(K)$ -space (Benyamini 1973).
- In the **complex case**: every **1-complemented** subspace of a space with 1-unconditional basis also has 1-unconditional basis. (Kalton-Wood 1976)

Recently (2021), **G. Plebanek** and **A. Salguero** have constructed an **example** (denoted by  $PS_2$ ) of a complemented subspace of  $C(K)$ -space which is not isomorphic to any  $C(K)$ -space.

## Conjectures (?)

- Every complemented subspace of  $L_1[0, 1]$  is isomorphic to  $\ell_1$  or  $L_1[0, 1]$ .
- Every complemented subspace of  $C[0, 1]$  is isomorphic to a  $C(K)$ -space.
- Every complemented subspace of a space with unconditional basis has unconditional basis.

# Some answers and some open questions

## Positive answers

- Every **1-complemented** subspace of an  $L_p$ -space ( $1 \leq p < \infty$ ) is an  $L_p$ -space (Bernau-Lacey 1974).
- Every **1-complemented** subspace of a **separable**  $C(K)$ -space is isomorphic to a  $C(K)$ -space (Benyamini 1973).
- In the **complex case**: every **1-complemented** subspace of a space with 1-unconditional basis also has 1-unconditional basis. (Kalton-Wood 1976)

Recently (2021), **G. Plebanek** and **A. Salguero** have constructed an **example** (denoted by  $PS_2$ ) of a complemented subspace of  $C(K)$ -space which is not isomorphic to any  $C(K)$ -space.

Actually,  $PS_2$  cannot be isomorphic to a Banach lattice.

## Conjectures (?)

- Every complemented subspace of  $L_1[0, 1]$  is isomorphic to  $\ell_1$  or  $L_1[0, 1]$ .
- Every complemented subspace of  $C[0, 1]$  is isomorphic to a  $C(K)$ -space.
- Every complemented subspace of a space with unconditional basis has unconditional basis.

# Table of Contents

- 1 A general overview of the question
- 2 Some remarks about  $PS_2$**
- 3 Concluding remarks

# $\mathcal{L}_p$ -spaces

**Definition.** Given  $1 \leq p \leq \infty$  and  $\lambda > 1$ , a Banach space  $X$  is an  $\mathcal{L}_{p,\lambda}$ -space if for every finite-dimensional subspace  $E$  of  $X$  there exists a finite-dimensional subspace  $F$  of  $X$  such that

- $E \subset F$ ;
- $d(F, \ell_p^{\dim F}) \leq \lambda$  (there is an isomorphism  $T : F \rightarrow \ell_p^{\dim F}$  such that  $\|T\| \|T^{-1}\| \leq \lambda$ ).

We say that a Banach space is an  $\mathcal{L}_p$ -space if it is an  $\mathcal{L}_{p,\lambda}$ -space for some  $\lambda > 1$ .



# $\mathcal{L}_p$ -spaces

**Definition.** Given  $1 \leq p \leq \infty$  and  $\lambda > 1$ , a Banach space  $X$  is an  $\mathcal{L}_{p,\lambda}$ -space if for every finite-dimensional subspace  $E$  of  $X$  there exists a finite-dimensional subspace  $F$  of  $X$  such that

- $E \subset F$ ;
- $d(F, \ell_p^{\dim F}) \leq \lambda$  (there is an isomorphism  $T : F \rightarrow \ell_p^{\dim F}$  such that  $\|T\| \|T^{-1}\| \leq \lambda$ ).

We say that a Banach space is an  $\mathcal{L}_p$ -space if it is an  $\mathcal{L}_{p,\lambda}$ -space for some  $\lambda > 1$ .

**Examples:**  $L_p(\mu)$ -spaces are  $\mathcal{L}_p$ -spaces for any  $1 \leq p \leq \infty$ .  $C(K)$ -spaces are  $\mathcal{L}_\infty$ -spaces.

# $\mathcal{L}_p$ -spaces

**Definition.** Given  $1 \leq p \leq \infty$  and  $\lambda > 1$ , a Banach space  $X$  is an  $\mathcal{L}_{p,\lambda}$ -space if for every finite-dimensional subspace  $E$  of  $X$  there exists a finite-dimensional subspace  $F$  of  $X$  such that

- $E \subset F$ ;
- $d(F, \ell_p^{\dim F}) \leq \lambda$  (there is an isomorphism  $T : F \rightarrow \ell_p^{\dim F}$  such that  $\|T\| \|T^{-1}\| \leq \lambda$ ).

We say that a Banach space is an  $\mathcal{L}_p$ -space if it is an  $\mathcal{L}_{p,\lambda}$ -space for some  $\lambda > 1$ .

**Examples:**  $L_p(\mu)$ -spaces are  $\mathcal{L}_p$ -spaces for any  $1 \leq p \leq \infty$ .  $C(K)$ -spaces are  $\mathcal{L}_\infty$ -spaces.

This property is preserved by **isomorphisms**.

# $\mathcal{L}_p$ -spaces

**Definition.** Given  $1 \leq p \leq \infty$  and  $\lambda > 1$ , a Banach space  $X$  is an  $\mathcal{L}_{p,\lambda}$ -space if for every finite-dimensional subspace  $E$  of  $X$  there exists a finite-dimensional subspace  $F$  of  $X$  such that

- $E \subset F$ ;
- $d(F, \ell_p^{\dim F}) \leq \lambda$  (there is an isomorphism  $T : F \rightarrow \ell_p^{\dim F}$  such that  $\|T\| \|T^{-1}\| \leq \lambda$ ).

We say that a Banach space is an  $\mathcal{L}_p$ -space if it is an  $\mathcal{L}_{p,\lambda}$ -space for some  $\lambda > 1$ .

**Examples:**  $L_p(\mu)$ -spaces are  $\mathcal{L}_p$ -spaces for any  $1 \leq p \leq \infty$ .  $C(K)$ -spaces are  $\mathcal{L}_\infty$ -spaces.

This property is preserved by **isomorphisms**.

Every **complemented subspace** of an  $\mathcal{L}_1$ -space (resp.,  $\mathcal{L}_\infty$ -space) is an  $\mathcal{L}_1$ -space (resp.,  $\mathcal{L}_\infty$ -space).

## $\mathcal{L}_p$ -spaces

**Definition.** Given  $1 \leq p \leq \infty$  and  $\lambda > 1$ , a Banach space  $X$  is an  $\mathcal{L}_{p,\lambda}$ -space if for every finite-dimensional subspace  $E$  of  $X$  there exists a finite-dimensional subspace  $F$  of  $X$  such that

- $E \subset F$ ;
- $d(F, \ell_p^{\dim F}) \leq \lambda$  (there is an isomorphism  $T : F \rightarrow \ell_p^{\dim F}$  such that  $\|T\| \|T^{-1}\| \leq \lambda$ ).

We say that a Banach space is an  $\mathcal{L}_p$ -space if it is an  $\mathcal{L}_{p,\lambda}$ -space for some  $\lambda > 1$ .

**Examples:**  $L_p(\mu)$ -spaces are  $\mathcal{L}_p$ -spaces for any  $1 \leq p \leq \infty$ .  $C(K)$ -spaces are  $\mathcal{L}_\infty$ -spaces.

This property is preserved by **isomorphisms**.

Every **complemented subspace** of an  $\mathcal{L}_1$ -space (resp.,  $\mathcal{L}_\infty$ -space) is an  $\mathcal{L}_1$ -space (resp.,  $\mathcal{L}_\infty$ -space).

**Definition.** A Banach lattice  $X$  is said to be an **AM-space** if  $\|x \vee y\| = \max\{\|x\|, \|y\|\}$  for any  $x, y \in X^+$ . An **AL-space** is a Banach lattice such that  $\|x + y\| = \|x\| + \|y\|$  for every  $x, y \in X^+$ .

## A first remark

$PS_2$  is complemented in a  $C(K)$ -space. Thus, it is an  $\mathcal{L}_\infty$ -space.

## A first remark

$PS_2$  is complemented in a  $C(K)$ -space. Thus, it is an  $\mathcal{L}_\infty$ -space.

**Consequence.** If  $PS_2$  were isomorphic to a Banach lattice, then it would be isomorphic to an AM-space. **Why?**

## A first remark

$PS_2$  is complemented in a  $C(K)$ -space. Thus, it is an  $\mathcal{L}_\infty$ -space.

**Consequence.** If  $PS_2$  were isomorphic to a Banach lattice, then it would be isomorphic to an AM-space. **Why?**

### Theorem

*Let  $X$  be a Banach lattice which is an  $\mathcal{L}_1$ -space. Then  $X$  is lattice isomorphic to an  $L_1(\mu)$ -space.*

## A first remark

$PS_2$  is complemented in a  $C(K)$ -space. Thus, it is an  $\mathcal{L}_\infty$ -space.

**Consequence.** If  $PS_2$  were isomorphic to a Banach lattice, then it would be isomorphic to an AM-space. **Why?**

### Theorem

Let  $X$  be a Banach lattice which is an  $\mathcal{L}_1$ -space. Then  $X$  is lattice isomorphic to an  $L_1(\mu)$ -space.

**Idea (Abramovich-Wojtaszczyk, 1975).** For every  $x \in X$ , define

$$\|x\| = \sup \left\{ \sum_{i=1}^m \|x_i\| : (x_i)_{i=1}^m \text{ with } |x_i| \wedge |x_j| = 0 \text{ s.t. } x = \sum_{i=1}^m x_i \right\}.$$



## A first remark

$\text{PS}_2$  is complemented in a  $C(K)$ -space. Thus, it is an  $\mathcal{L}_\infty$ -space.

**Consequence.** If  $\text{PS}_2$  were isomorphic to a Banach lattice, then it would be isomorphic to an AM-space. **Why?**

### Theorem

Let  $X$  be a Banach lattice which is an  $\mathcal{L}_1$ -space. Then  $X$  is lattice isomorphic to an  $L_1(\mu)$ -space.

**Idea (Abramovich-Wojtaszczyk, 1975).** For every  $x \in X$ , define

$$\|x\| = \sup \left\{ \sum_{i=1}^m \|x_i\| : (x_i)_{i=1}^m \text{ with } |x_i| \wedge |x_j| = 0 \text{ s.t. } x = \sum_{i=1}^m x_i \right\}.$$

$\|\cdot\|$  is an AL-norm (compatible with the lattice order of  $X$ ) and is related with the original norm by

$$\|x\| \leq \|x\| \leq (K_G \lambda)^2 \|x\|, \quad x \in X.$$

## Theorem

*Let  $X$  be a Banach lattice which is an  $\mathcal{L}_1$ -space. Then  $X$  is lattice isomorphic to an  $L_1(\mu)$ -space.*

## Theorem

Let  $X$  be a Banach lattice which is an  $\mathcal{L}_1$ -space. Then  $X$  is lattice isomorphic to an  $L_1(\mu)$ -space.

**Corollary 1.** Let  $X$  be a Banach lattice which is an  $\mathcal{L}_\infty$ -space. Then  $X$  is lattice isomorphic to an  $AM$ -space.

## Theorem

Let  $X$  be a Banach lattice which is an  $\mathcal{L}_1$ -space. Then  $X$  is lattice isomorphic to an  $L_1(\mu)$ -space.

**Corollary 1.** Let  $X$  be a Banach lattice which is an  $\mathcal{L}_\infty$ -space. Then  $X$  is lattice isomorphic to an  $AM$ -space.

**Proof.**  $X^*$  Banach lattice and  $\mathcal{L}_1$ -space, hence  $X^*$  is lattice isomorphic to an  $AL$ -space. Then,  $X^{**}$  is lattice isomorphic to certain  $C(K)$ -space, so  $X$  is lattice embeddable into that  $C(K)$ -space.  $\square$

## Theorem

Let  $X$  be a Banach lattice which is an  $\mathcal{L}_1$ -space. Then  $X$  is lattice isomorphic to an  $L_1(\mu)$ -space.

**Corollary 1.** Let  $X$  be a Banach lattice which is an  $\mathcal{L}_\infty$ -space. Then  $X$  is lattice isomorphic to an  $AM$ -space.

**Proof.**  $X^*$  Banach lattice and  $\mathcal{L}_1$ -space, hence  $X^*$  is lattice isomorphic to an  $AL$ -space. Then,  $X^{**}$  is lattice isomorphic to certain  $C(K)$ -space, so  $X$  is lattice embeddable into that  $C(K)$ -space.  $\square$

**Corollary 2.** If the CSP had a positive answer in the separable setting:

- ① Every complemented subspace of  $L_1[0, 1]$  would be isomorphic to  $\ell_1$  or to  $L_1[0, 1]$ .
- ② Every complemented subspace of  $C[0, 1]$  would be isomorphic to a  $C(K)$ -space.

## Some comments about $PS_2$

Let  $\mathcal{A} = \{A_\xi : \xi < \mathfrak{c}\} \subset \mathcal{P}(\mathbb{N})$  be an **almost disjoint family**, that is,  $|A_\xi|$  is infinite for every  $\xi$  and  $|A_\xi \cap A_{\xi'}|$  is finite whenever  $\xi \neq \xi'$ .

For every  $\xi < \mathfrak{c}$  we decompose  $A_\xi \times \{0, 1\} = \widehat{A}_\xi = B_\xi^0 \uplus B_\xi^1$  in the following way

## Some comments about $PS_2$

Let  $\mathcal{A} = \{A_\xi : \xi < \mathfrak{c}\} \subset \mathcal{P}(\mathbb{N})$  be an **almost disjoint family**, that is,  $|A_\xi|$  is infinite for every  $\xi$  and  $|A_\xi \cap A_{\xi'}|$  is finite whenever  $\xi \neq \xi'$ .

For every  $\xi < \mathfrak{c}$  we decompose  $A_\xi \times \{0, 1\} = \widehat{A}_\xi = B_\xi^0 \uplus B_\xi^1$  in the following way

$$\widehat{A}_\xi \left\{ \begin{array}{cccccccccccccccc} \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \dots \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \dots \end{array} \right.$$

## Some comments about $PS_2$

Let  $\mathcal{A} = \{A_\xi : \xi < \mathfrak{c}\} \subset \mathcal{P}(\mathbb{N})$  be an **almost disjoint family**, that is,  $|A_\xi|$  is infinite for every  $\xi$  and  $|A_\xi \cap A_{\xi'}|$  is finite whenever  $\xi \neq \xi'$ .

For every  $\xi < \mathfrak{c}$  we decompose  $A_\xi \times \{0, 1\} = \widehat{A}_\xi = B_\xi^0 \uplus B_\xi^1$  in the following way

$$\widehat{A}_\xi \left\{ \begin{array}{cccccccccccccccc} \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \dots \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \dots \end{array} \right.$$



## Some comments about $PS_2$

Let  $\mathcal{A} = \{A_\xi : \xi < \mathfrak{c}\} \subset \mathcal{P}(\mathbb{N})$  be an **almost disjoint family**, that is,  $|A_\xi|$  is infinite for every  $\xi$  and  $|A_\xi \cap A_{\xi'}|$  is finite whenever  $\xi \neq \xi'$ .

For every  $\xi < \mathfrak{c}$  we decompose  $A_\xi \times \{0, 1\} = \widehat{A}_\xi = B_\xi^0 \uplus B_\xi^1$  in the following way

$$\widehat{A}_\xi \left\{ \begin{array}{cccccccccccccccc} \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \dots \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \dots \end{array} \right.$$

We define:

$$JL(\mathcal{B}) = \overline{\text{span}}(\{\mathbf{1}_{B_\xi^0}, \mathbf{1}_{B_\xi^1} : \xi < \mathfrak{c}\} \cup c_{00}(\widehat{\mathbb{N}}) \cup \{\mathbf{1}_{\widehat{\mathbb{N}}}\}) \subset \ell_\infty,$$

$$JL(\mathcal{A}) = \overline{\text{span}}(\{\mathbf{1}_{\widehat{A}_\xi} : \xi < \mathfrak{c}\} \cup c_{00}(\widehat{\mathbb{N}}) \cup \{\mathbf{1}_{\widehat{\mathbb{N}}}\}) \subset \ell_\infty.$$

## Some comments about $PS_2$

Let  $\mathcal{A} = \{A_\xi : \xi < \mathfrak{c}\} \subset \mathcal{P}(\mathbb{N})$  be an **almost disjoint family**, that is,  $|A_\xi|$  is infinite for every  $\xi$  and  $|A_\xi \cap A_{\xi'}|$  is finite whenever  $\xi \neq \xi'$ .

For every  $\xi < \mathfrak{c}$  we decompose  $A_\xi \times \{0, 1\} = \widehat{A}_\xi = B_\xi^0 \uplus B_\xi^1$  in the following way

$$\widehat{A}_\xi \left\{ \begin{array}{cccccccccccccccc} \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \dots \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \dots \end{array} \right.$$

We define:

$$JL(\mathcal{B}) = \overline{\text{span}}(\{\mathbf{1}_{B_\xi^0}, \mathbf{1}_{B_\xi^1} : \xi < \mathfrak{c}\} \cup c_{00}(\widehat{\mathbb{N}}) \cup \{\mathbf{1}_{\widehat{\mathbb{N}}}\}) \subset \ell_\infty,$$

$$JL(\mathcal{A}) = \overline{\text{span}}(\{\mathbf{1}_{\widehat{A}_\xi} : \xi < \mathfrak{c}\} \cup c_{00}(\widehat{\mathbb{N}}) \cup \{\mathbf{1}_{\widehat{\mathbb{N}}}\}) \subset \ell_\infty.$$

These spaces can be identified with  $C(K)$ -spaces, with  $K$  scattered.

## Some comments about PS<sub>2</sub>

Let  $\mathcal{A} = \{A_\xi : \xi < \mathfrak{c}\} \subset \mathcal{P}(\mathbb{N})$  be an **almost disjoint family**, that is,  $|A_\xi|$  is infinite for every  $\xi$  and  $|A_\xi \cap A_{\xi'}|$  is finite whenever  $\xi \neq \xi'$ .

For every  $\xi < \mathfrak{c}$  we decompose  $A_\xi \times \{0, 1\} = \widehat{A}_\xi = B_\xi^0 \uplus B_\xi^1$  in the following way

$$\widehat{A}_\xi \left\{ \begin{array}{cccccccccccccccc} \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \dots \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \dots \end{array} \right.$$

We define:

$$\text{JL}(\mathcal{B}) = \overline{\text{span}}(\{\mathbf{1}_{B_\xi^0}, \mathbf{1}_{B_\xi^1} : \xi < \mathfrak{c}\} \cup c_{00}(\widehat{\mathbb{N}}) \cup \{\mathbf{1}_{\widehat{\mathbb{N}}}\}) \subset \ell_\infty,$$

$$\text{JL}(\mathcal{A}) = \overline{\text{span}}(\{\mathbf{1}_{\widehat{A}_\xi} : \xi < \mathfrak{c}\} \cup c_{00}(\widehat{\mathbb{N}}) \cup \{\mathbf{1}_{\widehat{\mathbb{N}}}\}) \subset \ell_\infty.$$

These spaces can be identified with  $C(K)$ -spaces, with  $K$  scattered.

Moreover, we can define a norm-one projection

$$P : \text{JL}(\mathcal{B}) \longrightarrow \text{JL}(\mathcal{A})$$

$$f \longmapsto Pf(n, 0) = Pf(n, 1) = \frac{f(n, 0) + f(n, 1)}{2}$$

We define  $X := \text{Ker}(P)$ , which is complemented in  $JL(\mathcal{B})$  by  $Q = \text{Id} - P$ .

We define  $X := \text{Ker}(P)$ , which is complemented in  $\text{JL}(\mathcal{B})$  by  $Q = \text{Id} - P$ .  
Since for every  $f \in \text{JL}(\mathcal{B})$

$$Qf(n, 0) = -Qf(n, 1) = \frac{f(n, 0) - f(n, 1)}{2},$$

then  $\|Q\| = 1$ .

We define  $X := \text{Ker}(P)$ , which is complemented in  $JL(\mathcal{B})$  by  $Q = \text{Id} - P$ .  
 Since for every  $f \in JL(\mathcal{B})$

$$Qf(n, 0) = -Qf(n, 1) = \frac{f(n, 0) - f(n, 1)}{2},$$

then  $\|Q\| = 1$ .

G. Plebanek and A. Salguero Alarcón show, through an inductive process of cardinality  $\aleph_1$ , that there exist almost disjoint families  $\mathcal{A}, \mathcal{B}$  such that  $X$  is not isomorphic to a  $C(K)$ -space. This  $X$  was christened  $PS_2$ .

# A simplification of our problem

Since  $PS_2 \subset \ell_\infty$ , then it has a **countable norming set**.

# A simplification of our problem

Since  $PS_2 \subset \ell_\infty$ , then it has a **countable norming set**.

Moreover,  $PS_2$  is 1-complemented in  $C(K)$ -space, with  $K$  scattered compact, so it is an **isometric predual of  $\ell_1(\Gamma)$** .



# A simplification of our problem

Since  $PS_2 \subset \ell_\infty$ , then it has a **countable norming set**.

Moreover,  $PS_2$  is 1-complemented in  $C(K)$ -space, with  $K$  scattered compact, so it is an **isometric predual of  $\ell_1(\Gamma)$** .

Consequently, the following statements are equivalent:

- 1  $PS_2$  is isomorphic to a Banach lattice.

# A simplification of our problem

Since  $PS_2 \subset \ell_\infty$ , then it has a **countable norming set**.

Moreover,  $PS_2$  is 1-complemented in  $C(K)$ -space, with  $K$  scattered compact, so it is an **isometric predual of  $\ell_1(\Gamma)$** .

Consequently, the following statements are equivalent:

- 1  $PS_2$  is isomorphic to a Banach lattice.
- 2  $PS_2$  is isomorphic to a sublattice of  $\ell_\infty$ .

# A simplification of our problem

Since  $PS_2 \subset \ell_\infty$ , then it has a **countable norming set**.

Moreover,  $PS_2$  is 1-complemented in  $C(K)$ -space, with  $K$  scattered compact, so it is an **isometric predual of  $\ell_1(\Gamma)$** .

Consequently, the following statements are equivalent:

- ①  $PS_2$  is isomorphic to a Banach lattice.
- ②  $PS_2$  is isomorphic to a sublattice of  $\ell_\infty$ .
- ③ There exists a norming sequence  $(x_n^*)_{n=0}^\infty$  in  $B_{PS_2^*}$  such that for every  $f \in PS_2$  there is an element  $g \in PS_2$  such that

$$x_n^*(g) = |x_n^*(f)|, \text{ for every } n \in \mathbb{N}.$$

# Table of Contents

- 1 A general overview of the question
- 2 Some remarks about  $PS_2$
- 3 Concluding remarks**

# Concluding remarks

**CSP in complex Banach lattices:** Is every complemented subspace of a complex Banach lattice isomorphic to a complex Banach lattice?

Recall that a **complex Banach lattice** is the complexification of a real Banach lattice  $X \oplus iX$  equipped with the norm  $\|x + iy\| := \||x + iy|\|_X$ , where

$$|x + iy| = \sup_{\theta \in [0, 2\pi]} \{x \cos \theta + y \sin \theta\}, \quad \text{for every } x, y \in X.$$

# Concluding remarks

**CSP in complex Banach lattices:** Is every complemented subspace of a complex Banach lattice isomorphic to a complex Banach lattice?

Recall that a **complex Banach lattice** is the complexification of a real Banach lattice  $X \oplus iX$  equipped with the norm  $\|x + iy\| := \||x + iy|\|_X$ , where

$$|x + iy| = \sup_{\theta \in [0, 2\pi]} \{x \cos \theta + y \sin \theta\}, \quad \text{for every } x, y \in X.$$

# Concluding remarks

**CSP in complex Banach lattices:** Is every complemented subspace of a complex Banach lattice isomorphic to a complex Banach lattice?

Recall that a **complex Banach lattice** is the complexification of a real Banach lattice  $X \oplus iX$  equipped with the norm  $\|x + iy\| := \||x + iy\|_X$ , where

$$|x + iy| = \sup_{\theta \in [0, 2\pi]} \{x \cos \theta + y \sin \theta\}, \quad \text{for every } x, y \in X.$$

- With **slight modifications**, it is possible to construct a  $\widetilde{\text{PS}}_2$  space such that  $\widetilde{\text{PS}}_2 \oplus i\widetilde{\text{PS}}_2$  is **not isomorphic to a complex Banach lattice**.

# Concluding remarks

**CSP in complex Banach lattices:** Is every complemented subspace of a complex Banach lattice isomorphic to a complex Banach lattice?

Recall that a **complex Banach lattice** is the complexification of a real Banach lattice  $X \oplus iX$  equipped with the norm  $\|x + iy\| := \||x + iy|\|_X$ , where

$$|x + iy| = \sup_{\theta \in [0, 2\pi]} \{x \cos \theta + y \sin \theta\}, \quad \text{for every } x, y \in X.$$

- With **slight modifications**, it is possible to construct a  $\widetilde{\text{PS}}_2$  space such that  $\widetilde{\text{PS}}_2 \oplus i\widetilde{\text{PS}}_2$  is **not isomorphic to a complex Banach lattice**.



# Concluding remarks

**CSP in complex Banach lattices:** Is every complemented subspace of a complex Banach lattice isomorphic to a complex Banach lattice?

Recall that a **complex Banach lattice** is the complexification of a real Banach lattice  $X \oplus iX$  equipped with the norm  $\|x + iy\| := \||x + iy|\|_X$ , where

$$|x + iy| = \sup_{\theta \in [0, 2\pi]} \{x \cos \theta + y \sin \theta\}, \quad \text{for every } x, y \in X.$$

- With **slight modifications**, it is possible to construct a  $\widetilde{\text{PS}}_2$  space such that  $\widetilde{\text{PS}}_2 \oplus i\widetilde{\text{PS}}_2$  is **not isomorphic to a complex Banach lattice**.
- The **separable case** of the CSP is still open!

# Thank you!